

SOIL MESOFAUNA

Soil fauna are commonly grouped together by taxonomy, trophic level, functional groups, and body size. These groupings, depicted in Figure 1, help describe who these fauna are and what they are doing in soils.

Soil mesofauna are tiny animals that range in size from 100µm to 2mm, which is comparable to a grain of sand. Prominent fauna in this group includes:

- Collembola (Springtails)
- Acari (Mites)
- Protura
- Diplura
- Enchytraidae (potworms)
- Pseudoscorpionida
- Myriapods

Mites and springtails are the most abundant mesofauna, examples of which are pictured in Figure 2. Soil mesofauna fill many diverse roles in soils due to their wide array of behaviors and habitat preferences. Trophic groupings describe what fauna are consuming. Functional groupings describe how fauna interact with their environment, and are influenced by how the fauna move through the soil which is determined by their body size.




FIGURE 1. Soil fauna groups defined by taxonomy, body width, and trophic group. The color of the shapes indicates the trophic groups; location of color is not related to size of fauna. The trophic groups are generalized for each taxonomic group.



FIGURE 2. (A) Collembola courtship display [family: Sminthuridae]. (B) Oribatid mite feeding on fungi [family: Euphthiracaridae]. (C) Collembola laying eggs [family: Isotomidae] (D) Mesostigmatid mite consuming a Collembola [family: Laelapidae]. Photography by Andy Murray: www.chaosofdelight.com



WHY DO WE CARE?

TROPHIC GROUPS

- Detritivores - consume plant litter and organic matter
- Microbivores - consume microbes
- Predators - consume other fauna
- Omnivores - consume a variety of flora and fauna

FUNCTIONAL GROUPS

- Nutrient Mineralizers - transform nutrients into plant available forms
- Litter Transformers - produce organic biogenic structures (fecal pellets)
- Ecosystem Engineers - transform soil into other structured materials

Soil mesofauna can fill a variety of different trophic groups and functional roles in soil ecosystems. These different groupings help us understand how soil fauna impact crop production.

Soil mesofauna are important because they impact many soil processes that affect crop production (Figure 3). Most notably, mesofauna stimulate soil organic matter decomposition and nutrient cycling. Mesofauna predominantly affect soil processes through their feeding habits, including:

- shredding plant litter and organic matter
- grazing on soil microbes (primarily fungi and bacteria)
- feeding on fungal seed coats and plant roots

In addition to having big impacts on nutrient dynamics in soils, through their behavior mesofauna can also influence:

- plant pathogen suppression and transmission
- seed germination
- root exudates
- plant nutrient allocation and growth

Though mesofauna primarily affect crop production through their interactions with other factors in the soil environment, these indirect effects can add up to big impacts on crops!




FIGURE 3. Visual overview of how soil mesofauna affect crop growth. Arrows depict drivers of soil processes.

FURTHER READINGS AND REFERENCES

Briones, M.J., 2014, Soil fauna and soil functions: a jigsaw puzzle. *Frontiers in Environmental Science*, Volume 2, 7pp.

Orgiazzi, A., Bardgett, R.D., Barrios, E., Behan-Pelletier, V., Briones, M.J.I., Chotte, J-L., De Deyn, G.B., Eggleton, P., Fierer, N., Fraser, T., Hedlund, K., Jeffery, S., Johnson, N.C., Jones, A., Kandeler, E., Kaneko, N., Lavelle, P., Lemanceau, P., Miko, L., Montanarella, L., Moreira, F.M.S., Ramirez, K.S., Scheu, S., Singh, B.K., Six, J., van der Putten, W.H., Wall, D.H. (Eds.), 2016, *Global Soil Biodiversity Atlas*. European Commission, Publications Office of the European Union, Luxembourg. 176 pp.



Created by Ashley Jernigan, Graduate Student, Department of Entomology,
Cornell University
Funded by Cornell Entomology Department Extension & Outreach Assistantship

Cornell
AgriTech
New York State Agricultural
Experiment Station

SOIL MESOFAUNA

Soil mesofauna can be found on every continent in the world in a variety of habitats. Soil mesofauna includes the fauna that live on the soil surface and in litter (epigeic) and those that reside lower within the soil profile (euedaphic). In agricultural soils, most biological activity occurs in the top 4 to 8 inches of soil.

Soil mesofauna populations vary greatly over time and space. Factors that affect the distribution and abundance of these fauna include: soil type, soil compaction, temperature, precipitation and moisture, and food resource availability. Soil mesofauna aggregate in favorable habitat conditions where they have the resources they need to survive, which causes very patchy distributions across land areas. In the Northeastern United States mesofauna abundances are greatest in the spring and fall due to milder temperatures and favorable soil moisture conditions.

There are a variety of different methods for collecting soil fauna that are typically based on their body size and mobility. Soil mesofauna can be collected using heat extraction methods that utilize Berlese-Tullgren funnels. This method relies on the mesofauna's natural behavior of moving deeper into the soil profile whenever conditions get too hot and dry.

WHERE ARE THEY?




FIGURE 1. (A) Diagram of basic components necessary to conduct a soil mesofauna extraction, with examples items for components. (B) Pictures of example modified funnel setups.



Evaluating the soil mesofauna present in a field requires three main steps: soil sample collection, mesofauna extraction, and sample processing. Each of these steps can be tailored based on an individual's goals, resources, and constraints. For example, you may want to see how a new crop management practice impacts the soil mesofauna or compare the soil mesofauna between different fields. The main thing to keep in mind is that uniformity in sample collection, extraction, and processing is necessary to be able to compare findings between samples.

After completing your first extraction it is important to practice sorting through samples to be able to find the small mesofauna. Tips for observing and counting the extracted mesofauna include:

- make sure you have good lighting
- try placing the sample on different color backgrounds (black or white)
- put the sample in a flat, shallow dish or container to improve clarity

STEPS TO DETERMINE THE MESOFAUNA IN YOUR SOILS:

1. Create a sampling plan and collect soil samples
 - Choose a time when conditions are favorable for soil mesofauna
 - Determine how many soil samples are necessary to capture the spatial variability in your field
 - Using a shovel, collect uniform soil samples to a depth of at least 4 inches
2. Extract mesofauna from soil samples
 - Set up modified Berlese-Tullgren funnel(s) following example diagram
 - Place soil samples on funnels and leave for uniform amount of time until extraction is complete, usually about three days
3. Processing the extracted soil mesofauna
 - Using a good magnifying glass or other magnification tool, aim for at least 30x magnification, observe the mesofauna extracted in each soil sample
 - Determine level of mesofauna identification you are comfortable with
 - Record observations of the number of mesofauna in each sample

It is easiest to begin with total mesofauna counts while you get comfortable with this process. Informative comparisons can be made with just measurements of total mesofauna abundance, though as you become more familiar with mesofauna identification you may choose to identify your fauna to broad taxonomic groups (total mites, total collembola, and other taxa) to get a better understanding of the communities.




FIGURE 2. (A) Image of extracted mesofauna sample. (B) Image of processed sample with soil removed.

FURTHER READINGS AND REFERENCES

Helmberger, M., Wickings, K. 2017. The Soil Animal Handbook, Version 2.1.1, Cornell University, Geneva, NY.

Menta, C., Conti, F.D., Pinto, S. Bodini, A. 2018. Soil Biological Quality index (QBS-ar): 15 years of application at global scale. Ecological Indicators. Volume 85, Pages 773-780.



Created by Ashley Jernigan, Graduate Student, Department of Entomology, Cornell University
Funded by Cornell Entomology Department Extension & Outreach Assistantship



SOIL MESOFAUNA

Soil mesofauna are affected by environmental factors including soil type, structure, temperature, moisture, and food resource availability. Crop management decisions impact many of these factors in manners that either benefit or harm the mesofauna. Soil mesofauna communities can be highly diverse, and therefore contain many different species that respond differently to environmental conditions. However, generalization can be made about how environmental conditions and crop management practices impact the overall activity and functioning of soil mesofauna communities (Table 1).

TABLE 1. Generalized overview of how environmental conditions are impacted by crop management practices and how those changes affect the overall activity and functioning of soil mesofauna communities.

CROP MANAGEMENT PRACTICE	ENVIRONMENTAL CONDITION	EFFECT ON MESOFAUNA ACTIVITY
Crop Rotation	Food Resources	Positive
Cover Cropping	Soil Structure Soil Temperature Soil Moisture Food Resources	Positive
Intercropping	Food Resources Soil Structure	Positive
Organic Matter Inputs	Food Resources	Positive
Tillage	Soil Structure	Negative
Irrigation	Soil Moisture Soil Temperature Food Resources	Positive
Pesticides	Food Resources	Variable

When trying to manage your fields to enhance soil biological activity it is important to focus on creating a habitat that can support a diverse and active soil mesofauna community. The responsiveness of different species within the mesofauna community to changing environmental conditions leads to the high spatial variability we observe with mesofauna.



WHAT AFFECTS THEM?

In October 2020, multiple farms across the Finger Lakes Region of New York State which varied greatly in crop rotation and management history were sampled to assess their mesofauna communities. Across all of these samples, the total number of soil mesofauna found in a single sample ranged from 58 – 1678 individuals per kilogram (approximately 2 lbs) of dry soil, with a average of 545 individuals per kilogram of dry soil. The average mesofauna abundance observed is depicted in Figure 1.

The different farms and fields sampled varied greatly in mesofauna abundance due to the varying environmental conditions. Crop diversity, both incorporating more diverse crop species in rotations and increasing diversity through intercropping practices, had positive effects on mesofauna abundance and diversity. The rough amount of ground cover was also related to the mesofauna abundance, with more dense ground cover promoting a more abundant mesofauna community. The application of insecticides to control crop pests decreased the amount of mesofauna. Overall, increasing the number of management decisions which reduce soil disturbance and feeds the soil's biological food web leads to more abundant and diverse mesofauna communities.




FIGURE 1. Bar graph illustrating the high variability of the average mesofauna abundances found on farms across the Finger Lakes Region of New York State.

FURTHER READINGS AND REFERENCES

Moebius-Clune, B.N., D.J. Moebius-Clune, B.K. Gugino, O.J. Idowu, R.R. Schindelbeck, A.J. Ristow, H.M. van Es, J.E. Thies, H.A. Shayler, M.B. McBride, K.S.M Kurtz, D.W. Wolfe, and G.S. Abawi, 2016. Comprehensive Assessment of Soil Health – The Cornell Framework, Edition 3.2, Cornell University, Geneva, NY.

Bedano, J.C., Domínguez, A., Arolfa, R., Wall, L.G. 2016. Effect of Good Agricultural Practices under no-till on litter and soil invertebrates in areas with different soil types. Soil and Tillage Research. Volume 158, Pages 100-109.



Created by Ashley Jernigan, Graduate Student, Department of Entomology,
Cornell University
Funded by Cornell Entomology Department Extension & Outreach Assistantship

Cornell
AgriTech
New York State Agricultural
Experiment Station